

Vietnam Journal of Chemistry, International Edition, **55**(5): 569-572, 2017
DOI: 10.15625/2525-2321.2017-00509

Some triterpenoids and steroids from *Bruguiera cylindrica* leaves collected from Can Gio mangrove forest

Nguyen Thi Huong Que¹, Tran Minh Chi², Pham Nguyen Kim Tuyen², Nguyen Kim Phi Phung^{1*}

¹ University of Science, National University Ho Chi Minh City, Vietnam

² Sai Gon University, Ho Chi Minh City, Vietnam

Received 3 March 2017; Accepted for publication 20 October 2017

Abstract

Bruguiera cylindrica (Rhizophoraceae), distributed in the Southeast Asia, has been used as a traditional medicine in the treatment of diarrhea and healing of wounds. Some triterpenoids and steroids were isolated from the dried leaves of *Bruguiera cylindrica*, collected at Can Gio mangrove forest, Ho Chi Minh City, including lupeol (**1**), 3 β -hydroxyoleana-9(11),12-diene (**2**), a mixture of stigmasterol (**3a**) and β -sitosterol (**3b**), cholesta-4-ene-3-one (**4**) and 3 β -hydroxycholesta-5-ene-7-one (**5**). Their chemical structures were elucidated by spectroscopic analysis as well as comparing their data with the ones in the literature. Although these compounds were known in other plants this is the first time they are reported in this species.

Keywords. *Bruguiera cylindrica*, triterpenoids, steroids.

1. INTRODUCTION

Bruguiera cylindrica Blume (Rhizophoraceae), a mangrove plant, is distributed widely in Southeast Asia [6]. This plant has been used in Vietnamese traditional medicine for the treatment of diarrhea and healing of wounds [6, 7]. Four species of this genus are found in Vietnam including *B. cylindrica*, *B. gymnorhiza*, *B. parviflora* and *B. sexangula* [1]. In the study on Vietnamese mangrove plants, we investigated the chemical constituents of *B. cylindrica* leaves collected in Can Gio mangrove forest, Ho Chi Minh City, Viet Nam.

2. EXPERIMENTAL

Leaves of *Bruguiera cylindrica* were collected at Can Gio mangrove forest in Ho Chi Minh city, Vietnam in September 2015. The scientific name of the plant was authenticated by Dr. Pham Van Ngot, Faculty of Biology, Ho Chi Minh city Pedagogical University. A voucher specimen (**No US-B020**) was deposited in the herbarium of the Department of Organic Chemistry, University of Science, National University - Ho Chi Minh City.

Column chromatography was carried out on silica gel 60 F₂₅₄ (Merck) and gel Sephadex LH-20.

Extraction of Bruguiera cylindrica

The air-dried leaves of *Bruguiera cylindrica* (6

kg) were dried, ground then extracted with *n*-hexane, ethyl acetate, and methanol (3 x 20 L for each solvent), successively. The *n*-hexane extract (25 g) was subjected to column chromatography, eluted with a gradient of *n*-hexane/acetone to afford nine fractions (A1–A9). Fraction A3 (1.5 g) was purified by silica gel column chromatography to afford compound **1** (10 mg). Compound **3** (5 mg) was isolated from the fraction A5. Fraction A7 (2.49 g) was applied to silica gel column chromatography to give compound **4** (5 mg) and **5** (4 mg).

The ethyl acetate extract (100 g) was subjected to silica gel column chromatography, eluted with a gradient of *n*-hexane/ethyl acetate to afford nine fractions (B1–B9). Fraction B4 was subjected to Sephadex LH-20 column to give five fractions (B4.1–B4.5). Fraction B4.4 was further purified by silica gel column chromatography (chloroform/methanol, 95.5/0.5) to give compound **2** (5 mg).

3. RESULTS AND DISCUSSION

Compound **1** obtained as a white crystal is a triterpene due to the positive sulfuric acid test. The ¹H-NMR spectral data of compound **1** gave signals of an isopropenyl group at δ 4.72 (1H, *d*, *J* = 2.0 Hz, H-29a), 4.61 (1H, *dd*, *J* = 2.0, 1.5 Hz, H-29b) and 1.72 (3H, *s*, H-30), a methine proton at δ 3.16 (1H, *dd*, *J* = 11.5, 4.5 Hz, H-3), six singlets belonging to six

methyl groups, including δ 0.79 (3H, *s*, H-24), 0.86 (3H, *s*, H-28), 0.91 (3H, *s*, H-25), 0.99 (3H, *s*, H-23), 1.02 (3H, *s*, H-27) and 1.11 (3H, *s*, H-26). The ^{13}C -

NMR spectral data of compound **1** are shown in table 1. All the spectral data of compound **1** were similar to the one of lupeol in the literature [2, 5].




Figure 1: Chemical structure of isolated compounds

Compound **2** obtained as a white crystal. The molecular formula was established as $\text{C}_{30}\text{H}_{48}\text{O}$ from the APCI-MS data at m/z 425.44 $[\text{M}+\text{H}]^+$ (calc. 425.37). The ^1H -NMR spectral data of compound **2** gave one oxymethine doublet of doublet at δ 3.24 (1H, *dd*, $J = 12.0, 5.0$ Hz, H-3), two signals of olefin protons at δ 5.57 (1H, *d*, $J = 5.5$ Hz, H-11) and 5.50 (1H, *d*, $J = 5.5$ Hz, H-12), eight singlets belonging to six methyl groups, including δ 0.81 (3H, *s*, H-24), 0.87 (3H, *s*, H-23), 0.89 (3H, *s*, H-29), 0.90 (3H, *s*, H-30), 0.99 (3H, *s*, H-28), 1.03 (3H, *s*, H-27), 1.14 (3H, *s*, H-25) and 1.19 (3H, *s*, H-26). The ^{13}C -NMR spectral data of compound **2** are shown in table 1. All the spectral data of compound **2** were similar to the ones of 3β-hydroxyoleana-9(11),12-diene in the

literature [4, 5].

Compound **3** (**3a** and **3b**) was isolated as a white amorphous powder. The APCI-MS gave a molecular ion peaks at m/z 397.36 $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$ (calc. 397.38) of β-sitosterol and 395.42 $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$ (calc. 395.38) of stigmasterol, the molecular ion peak of β-sitosterol is taller. The ^1H -NMR spectrum of **3** showed signals of olefin protons at δ 5.35 (3H, *d*, $J = 5.5$ Hz, H-6, H-6'), 5.15 (1H, *dd*, $J = 15.0, 8.5$ Hz, H-22) and 5.01 (1H, *dd*, $J = 15.0, 8.5$ Hz, H-23), along with one oxymethine multiplet at δ 3.52 (3H, *m*, H-3, H-3') and many other signals in the high field from δ 2.5-1.0. The ^{13}C -NMR spectrum showed four olefin carbon signals at δ 140.9 (C-5, C-5'), three olefin methines

at δ 138.5 (C-22), 129.5 (C-23), 121.9 (C-6, C-6') and one oxygenated methine at δ_C 71.9 (C-3, C-3') of a mixture β -sitosterol and stigmasterol [3]. All

signals showed that **3** was a mixture of β -sitosterol and stigmasterol with the ratio of 1:2 based on their protons integrations.

Table 1: NMR data of compounds **1**, **2**, **4** and **5**

Po s.	Compound 1 (CD ₃ OD)		Compound 2 (CDCl ₃)		Compound 4 (CDCl ₃)		Compound 5 (CDCl ₃)	
	δ_C	δ_H, J (Hz)	δ_C	δ_H, J (Hz)	δ_C	δ_H, J (Hz)	δ_C	δ_H, J (Hz)
1	38.8		38.7		35.9		36.5	
2	27.4		27.9		34.1		31.4	
3	79.0	3.16 (1H, <i>dd</i> , 11.5, 4.5)	78.7	3.24 (1H, <i>dd</i> , 12.0, 5.0)	199.7		70.7	3.67 (1H, <i>m</i>)
4	38.7		38.9		123.8	5.72 (1H, <i>s</i>)	38.9	
5	55.4		51.2		171.8		165.2	
6	18.3		18.4		33.1		126.3	5.69 (1H, <i>s</i>)
7	34.3		32.1		32.2		202.5	
8	40.9		37.1		35.8		45.6	
9	50.5		154.3		54.0		50.1	
10	37.2		40.7		38.8		38.4	
11	21.0		115.7	5.57 (1H, <i>d</i> , 5.5)	21.2		21.4	
12	25.2		120.7	5.50 (1H, <i>d</i> , 5.5)	39.8		42.9	
13	38.1		147.1		42.6		43.3	
14	42.9		42.8		56.0		50.2	
15	27.5		25.7		24.3		26.3	
16	35.6		27.2		28.3		28.7	
17	43.0		31.9		56.2		54.9	
18	48.4		45.6		12.1	0.71 (3H, <i>s</i>)	12.1	0.68 (3H, <i>s</i>)
19	48.0	2.44 (1H, <i>m</i>)	46.9		17.5	1.17 (3H, <i>s</i>)	17.5	1.20 (3H, <i>s</i>)
20	151.0		31.1		35.8		35.9	
21	29.9		34.6		18.9	0.91 (3H, <i>d</i> , 6.5)	19.1	0.93 (3H, <i>d</i> , 6.5)
22	40.0		37.2		36.3		36.2	
23	27.9	0.99 (3H, <i>s</i>)	28.2	0.87 (3H, <i>s</i>)	23.2		23.2	
24	15.4	0.79 (3H, <i>s</i>)	15.4	0.81 (3H, <i>s</i>)	39.7		40.0	
25	16.1	0.91 (3H, <i>s</i>)	20.1	1.14 (3H, <i>s</i>)	28.3		29.3	
26	16.0	1.11 (3H, <i>s</i>)	21.0	1.19 (3H, <i>s</i>)	22.8	0.84 (3H, <i>d</i> , 7.0)	23.2	0.84 (3H, <i>d</i> , 7.0)
27	14.6	1.02 (3H, <i>s</i>)	25.3	1.03 (3H, <i>s</i>)	22.9	0.81 (3H, <i>d</i> , 7.0)	21.6	0.82 (3H, <i>d</i> , 7.0)
28	18.0	0.86 (3H, <i>s</i>)	28.7	0.99 (3H, <i>s</i>)				
29	109.3	4.72 (1H, <i>d</i> , 2.0) 4.61 (1H, <i>dd</i> , 2.0, 1.5)	23.7	0.89 (3H, <i>s</i>)				
30	19.3	1.72 (3H, <i>s</i>)	33.2	0.90 (3H, <i>s</i>)				

Compound **4** was isolated as a white amorphous powder. Based on the HR-ESI-MS spectrum, the molecular formula of **4** was determined as C₂₇H₄₄O with a pseudomolecular ion peak at m/z 385.3448 [M+H]⁺ (calc. for C₂₇H₄₄O+H, 385.3470). The ¹H-NMR spectrum of **4** displayed a singlet at δ 5.72 (1H, *s*, H-4), three methyl doublets

at δ 0.91 (3H, *d*, J = 6.5 Hz, H-21), 0.84 (3H, *d*, J = 7.0 Hz, H-26), 0.81 (3H, *d*, J = 7.0 Hz, H-27), and two methyl singlets at δ 0.71 (3H, *s*, H-18), and 1.17 (3H, *s*, H-19). The ¹³C-NMR spectral data of compound **4** were shown in table 1. All signals showed that **4** was cholesta-4-ene-3-one due to the compatibility of its spectroscopic data with the

one in the literature [8].

Compound **5** was isolated as a white amorphous powder. The ^1H -NMR spectrum of **5** showed one oxymethine multiplet at δ 3.67 (1H, *m*, H-3), an olefin singlet at δ 5.69 (1H, *s*, H-6), three methyl doublets at δ 0.93 (3H, *d*, $J = 6.5$ Hz, H-21), 0.84 (3H, *d*, $J = 7.0$ Hz, H-26), 0.82 (3H, *d*, $J = 7.0$ Hz, H-27), and two methyl singlets at δ 0.68 (3H, *s*, H-18) and 1.20 (3H, *s*, H-19). The ^{13}C -NMR spectral data of compound **5** were shown in table 1. All signals showed that **5** was 3β -hydroxycholesta-5-ene-7-one due to the compatibility of its spectroscopic data with the one in the literature [9].

REFERENCES

1. Phạm Hoàng Hộ. *Cây cỏ Việt Nam*, Youth Publishing House, **2**, 111-120 (2003).
2. Darcy Burns, William F. Reynolds, Greg Buchanan, Paul B. Reese and Raul G. Enriquez. *Assignment of ^1H and ^{13}C -NMR spectra and investigation of hindered side-chain rotation in lupeol derivatives*, Magn. Reson. Chem., **38**, 488-493 (2000).
3. M. Rowshanul Habib, Farjana Nikkon, Matiar Rahman, M. Ekramul Haque and M. Rezaul Karim, *Isolation of stigmastrol and β -sitosterol from methanolic extract of root bark of Calotropis gigantea (Linn.)*, Pakistan Journal of Biological Science, **10(22)**, 4174-4176 (2007).
4. Reiko Tanaka and Shunyo Matsunaga. *Triterpene dienols and other constituents from the bark of Phyllanthus flexuosus*, Phytochemistry, **27(7)**, 2213-2217 (1988).
5. Shashi B. Mahato and Asish P. Kundu. *^{13}C -NMR spectra of pentacyclic triterpenoids. A compilation and some salient features*, Phytochemistry, **37(6)**, 1548 (1994).
6. Suchada Chantrapromma, Hoong-Kun Fun, Ibrahim Abdul Razak, Surat Laphookhieo and Chatchanok Karalai. *Absolute configuration of 3 α -feruloyltaraxerol dichloromethane solvate*, Acta. Cryst., **59**, 1864-1866 (2003).
7. Sundaram Ravikumar, Samuel Jacob Inbaneson, Palavesam Suganthi, Murugesan Gnanadesigan. *In vitro antiplasmodial activity of ethanolic extracts of mangrove plants from South East coast of India against chloroquine-sensitive Plasmodium falciparum*, Parasitol Res., **108**, 873-878 (2011).
8. Youngsheng Wang, Lili Wang, Yan Hu, Lantong Zhang and Zhihan Wang. *Isolation and identification of two steroid compounds from Ovidustus ranee*, Natural Product Research, **24(16)**, 1518-1522 (2010).
9. Yuchang Liu, Guiying Chen, Fanglan Ge, Wei Li, Lihua Zeng and Weiguo Cao. *Efficient biotransformation of cholesterol to androsta-1,4-diene-3,17-dione by a newly isolated actinomycete Gordonia neofelifaecis*, World J. Microbiol. Biotechnol., **27**, 759-765 (2011).

Corresponding author: **Nguyen Kim Phi Phung**

Department of Chemistry, University of Science, National University – HCM City
227, Nguyen Van Cu Street, District 5, Ho Chi Minh city, Vietnam
E-mail: kimphiphung@yahoo.fr; Telephone: 01226966660.